

Biomass, chlorophyll and nitrogen content of leaves of two chili pepper varieties (*Capsicum annuum*) in different fertilization treatments

SUHARJA^{1,*}, SUTARNO²

¹ SMA Negeri 1 Klaten. Jl. Merbabu 13, Gayampit, Klaten Selatan, Klaten 57423, Central Java, Indonesia. Tel./Fax.: +62-272-321150

² Bioscience Program, School of Graduates, Sebelas Maret University, Surakarta 57126, Central Java, Indonesia

Manuscript received: 28 October 2008. Revision accepted: 27 January 2009.

Abstract. *Suharja, Sutarno. 2009. Biomass, chlorophyll and nitrogen content of leaves of two chili pepper varieties (Capsicum annuum) in different fertilization treatments. Nusantara Bioscience 1: 9-16.* This study aims to determine the influence of various fertilization treatments on biomass, chlorophyll and nitrogen content of leaves from two varieties of chili, *Sakti* (large chili) and *Fantastic* (curly chili). The study was conducted in the village of Gatak, Karangnongko sub-district, Klaten District, Central Java in September 2006 to March 2007. The study used a complete block design with two factorial of chili varieties and fertilizer treatment. Fertilization treatments includes no fertilizer (control) (P1); manure 2 kg/plant (P2), manure (1 kg/plant) + chemical fertilizer (ZA, SP-36, KCl = 2: 1: 1) + NPK (P3); and manure (1 kg/plant) + chemical fertilizer (SP-36: KCl = 1:1) + liquid organic fertilizer (P4). Chlorophyll content was measured refers to Harborne (1987), whereas leaf nitrogen concentration was measured with Kjeldahl method. Data were analyzed using ANOVA followed by DMRT. The results showed that on the *Fantastic* chili fertilizer treatment affected the biomass and chlorophyll *a*, but gave no effect on chlorophyll *b*, total chlorophyll and leaf nitrogen. On the curly chili fertilizer treatment effected plant fresh weight, chlorophyll *a* and total chlorophyll, but gave no effect on dry weight, fresh fruit weight, chlorophyll *b* and leaf nitrogen. It is, therefore, recommended to use the formulation of manure + chemical fertilizer (SP-36: KCl = 1: 1) + liquid organic fertilizer in the cultivation of chili.

Key words: biomass, chlorophyll, leaf nitrogen, chili, *Capsicum annuum*, fertilizing.

Abstrak. *Suharja, Sutarno. 2009. Biomassa, kandungan klorofil dan nitrogen daun dua varietas cabai (Capsicum annuum) pada berbagai perlakuan pemupukan. Nusantara Bioscience 1: 9-16.* Penelitian ini bertujuan untuk mengetahui pengaruh berbagai perlakuan pemupukan terhadap biomassa, kandungan klorofil dan nitrogen daun dari dua varietas cabai, *Sakti* (cabai besar) dan *Fantastic* (cabai keriting). Penelitian dilakukan di Desa Gatak, Kecamatan Karangnongko, Kabupaten Klaten, Jawa Tengah pada September 2006 sampai Maret 2007. Penelitian menggunakan rancangan blok lengkap dengan dua faktorial yaitu varietas cabai dan perlakuan pemupukan. Perlakuan pemupukan meliputi tanpa pupuk (kontrol) (P1); pupuk kandang 2 kg/tanaman (P2), pupuk kandang (1 kg/tanaman) + pupuk kimia (ZA, SP-36, KCl = 2: 1: 1) + NPK (P3); dan pupuk kandang (1 kg/tanaman) + pupuk kimia (SP-36: KCl = 1:1) + pupuk organik cair (P4). Kadar klorofil diukur merujuk Harborne (1987), sedangkan kadar nitrogen daun diukur dengan metode Kjeldahl. Data dianalisis menggunakan Analisis Varians dilanjutkan DMRT. Hasil penelitian menunjukkan pada cabai *Fantastic*, perlakuan berbagai macam pemupukan berpengaruh terhadap biomassa dan klorofil *a*, namun tidak berpengaruh terhadap kandungan klorofil *b*, total klorofil dan nitrogen daun. Pada cabai *Sakti* perlakuan pemupukan berpengaruh terhadap bobot segar tanaman, kandungan klorofil *a* dan total klorofil, namun tidak berpengaruh terhadap bobot kering, bobot buah segar, kandungan klorofil *b* dan nitrogen daun. Oleh karena itu direkomendasikan untuk menggunakan formulasi pupuk kandang + pupuk kimia (SP-36: KCl = 1: 1) + pupuk organik cair dalam budidaya cabai.

Kata kunci: biomassa, klorofil, nitrogen daun, cabai, *Capsicum annuum*, pemupukan.

INTRODUCTION

The demand of chili pepper (*Capsicum annum* L) always increases with the increase of the number of food factory, family needs and various instant noodles, sauce, and chilly industry. Results of the 2007 chili crop was 6.30 tons/ha, which was lower than in 2006 6.51 tons/ha. Meanwhile import of pepper in 2006 was 11,885,501 tons and the export of pepper in 2006 was 8,004,450 tons. This condition implies that the production of large chili per hectare needs to be developed. Therefore it is necessary to find cultivation technology which can increase growth and yield of chili (www.hortikultura.deptan.go.id 2008).

According to Nyakpa et al. (1988) the success of a farming business is largely determined by the growth and yield of cultivated plants. If the growth and yield is satisfactory then the business is said to be successful. Allabi (2005), further states that peppers will give good results if the essential elements needs are met. To meet the needs of the essential elements, fertilization can be done. The successful use of organic fertilizers in encouraging crop production is not doubted. The provision of organic fertilizer to increase growth and yield can give better yield than that of phosphorus.

Meanwhile Sadewa (2008) states that the type of fertilizer mixture (N: P: K = 8.31 g: 12.21 g: 8.81 g) can improve the growth of plant's height, root's length and

chlorophyll content of three varieties of chili peppers at the vegetative phase. According to Chellemi and Lazarovits (2002) applying organic fertilizer 310 and 400 kg N/ha may cause a decrease in production of chili peppers and tomatoes, even providing 560 kg N/ha may lead to increase soil pH, NH₃ and the total number of fungus/mushroom, which can be toxic and can cause death of the pepper.

Based on the researches of Sadewo (2008), Allabi (2005) and Chellemi and Lazarovits (2002), chili pepper plants require macro and micro nutrients for growing and increasing the production. These needs of nutrients can be met through the provision of organic fertilizer and inorganic fertilizer. Formulation of the appropriate types of fertilizer can affect the biomass, chlorophyll content and the nitrogen of the leaf. The thought that says organic fertilizers can increase agricultural production is not entirely correct. Only the appropriate formulations of fertilizer that can affect the growth and the yield of chili. Therefore, research to find a formulation of organic and inorganic fertilizers and/or the combination of both fertilizers and its influence on the pepper plant needs to be done.

Big or large chili (*C. annuum*) has many varieties, among others: the long red chili (*C. annuum* var. *longum* (DC.) Sendt), round peppers (*C. annuum* var. *Cerasiforme* (Miller) Irish), sweet chili or bell peppers or paprika (*C. annuum* var. *grossum*) and green peppers (*C. annuum* var. *annuum*) and others. Curly chili is one of the varieties of long red chili (*C. annuum* var. *longum* (DC.) Sendt) (Pracaya 2000; Setiadi 1993).

Biomass is defined as the sum total of life at any given time and a certain area. Biomass can be expressed as the biomass volume, wet weight biomass, dry weight biomass and organo biomass (Michael 1994). Prawirohatmodjo *et al.* (2001) further state that biomass covers the entire body of a living creature, even when the body is only a branch or a leaf of a tree as long as still attached to the plants. According to Salisbury and Ross (1995) fresh mass is determined by harvesting the whole plants or parts of the plant and weigh them quickly before too much water evaporates from the material. About 75% of plant biomass is produced several weeks before harvest time so that at that time the plant needs higher nutrient and it absorbs fertilizer more efficiently (Rubatzky and Yamaguchi 1995). The measurement of biomass can also use dry mass. Dry mass measurement needs to be done, because of various problems arising from the content of the water so that the productivity of cultivated plants should also be measured by using the dry mass of the plants (Salisbury and Ross 1995). Dry weight biomass is measured to obtain the overall appearance of plant growth (Sitompul and Guritno 1995). Measurement of dry weight accumulation is an analogy to know the distribution pattern of assimilation from the source to the target (Gardner *et al.* 1991).

Chlorophyll is a magnesium-porphyrin attached to proteins (Nelson and Cox 2004). Chlorophyll is an important catalyst for photosynthesis found in thylakoid membranes as a green pigment in plant photosynthetic tissues, which is loosely bound to proteins but easily extracted into a solvent such as acetone and ether lipids (Harborne 1987). Taller plants have two types of

chlorophyll that are chlorophyll *a* and chlorophyll *b*. Chlorophyll *a* is a complex compound of magnesium and porphyrin which has cyclopentanone ring (ring V). The four nitrogen atoms linked by ties of coordination with Mg²⁺ forming a firm planar compound. The hydrophobic side chain which is a terpenoid alcohol, or phytol, which are connected by ester bonds propionate group of ring IV. Chlorophyll *b* is the second chlorophyll found in plants (Wirahadikusumah 1985). The structure of chlorophyll *b* is different from chlorophyll *a* because chlorophyll *a* has a methyl catalyst, while chlorophyll *b* has an aldehyde group which is attached on the right top of the pyrrole ring (Harborne 1987). Chlorophyll *b* may derive from chlorophyll *a*, methyl groups are oxidized on its second ring and become the aldehyde group, or are possible for the porphyrin compounds which can be converted to chlorophyll *a* and *b* (Bonner and Varner 1965). Porphyrin in chlorophyll *a* is the precursor of chlorophyll both *a* and *b* (Mohr and Schopfer 1995).

In plant tissues, nitrogen is a constituent component of many essential compounds such as proteins, amino acids, amides, nucleic acids, nucleotides, coenzymes (Loveless 1987), chlorophyll, cytosine, auxin (Lakitan 2007), and the main components of dry material derived from protoplasmic material plants (Salisbury and Ross 1992). Plants absorb nitrogen element in the form of NO³⁻ and NH⁴⁺ (Nyakpa *et al.* 1988).

Fertilizers are all materials provided for the ground to improve the physical, chemical and biological soil condition (Subagyo 1970). Fertilizer is a material that is given to soil, both organic and inorganic material, with objective to replace the loss of nutrients from the soil and to increase crop production (Sutejo 2002). Provision of different types of fertilizer can affect the growth and yield of plants. Yield and quality of paprika will be different depends on the different types of nitrogen fertilizer which is given to it, both PCU (*polyolefinresin coated urea*) and SCU (*sulfur coated urea*) (Guertal 2000).

Organic fertilizers have a role in influencing the physical properties, chemical and soil biological activity. Organic fertilizers can improve soil physical characters through the formation of soil aggregate structure and a stable and closely related to water binding ability, water infiltration, reduce erosion, increase ion exchange capacity (CEC) and as a regulator of soil temperature, all affect the plant growth (Kononova 1999; Foth 1990). Organic fertilizers contain nutrients that are needed by plant growth (Rauf 1995; Tandisau and Sariubang 1995). The use of organic extract (organic liquid fertilizer) with a concentration of 2-3 mL/L water can increase the yield of various crops, like peppers, tomatoes, and corn by about 25% (Sima 1999, 2005). Giving the organic extract with relatively short interval (7 days), can directly maintain the supply of nutrients and soil microbe vitamins that play a role in the decomposition process of soil organic matter and keep soil health (Diver 2001; Scheuerell 2004).

This study aims to know the influence of various fertilization treatments on biomass, chlorophyll content and leaf nitrogen in two varieties of chili pepper (*Capsicum annuum* L.), large chili and curly chili.

MATERIALS AND METHODS

Time and place

This research was conducted from September 2006 until March 2007 in the rice field in Gathak village, Karangnongko sub-district, Klaten District, Central Java. The measurement of chlorophyll content and leaf nitrogen, was conducted at the Laboratory of Soil Science, Faculty of Agriculture, Sebelas Maret University, Surakarta.

Material

Seeds of large chili from varieties of *Fantastic* and seed of curly chili varieties of *Sakti* which are from the same broodstock. It needs fertilizers such as NPK *Mutiara*, cattle manure, ZA, SP-36, KCl, and liquid organic fertilizer branded *Batari Sri*. According to PT. Batari Sri (2005), liquid organic fertilizer (LOF) *Batari Sri* is pure organic fertilizers that 97% are made from cattle urine and 3% of natural ingredients processed by fermentation to produce liquid fertilizer that do not contain zinc (Zn), copper (Cu) and lead (Pb).

Study design

This research was a factorial with randomized complete block design (RCBD) with 2 factors: (i) variety (two levels) (ii) fertilization (four levels), with 3 replications and each replication consisted of 20 planted chili. Fertilization treatments included: P1 = no fertilizer as control; P2 = manure 2 kg/plant; P3 = manure (1 kg/plant) + chemical fertilizer (ZA, SP-36, KCl = 2: 1: 1) + NPK *Mutiara*; and P4 = manure (1 kg/plant) + chemical fertilizer (SP-36: KCl = 1:1) + liquid organic fertilizer *Bathari Sri*.

Research parameter

The parameters of this study are: biomass (plant fresh weight, dry weight of plants, and fruit fresh weight per plant), chlorophyll (chlorophyll *a*, chlorophyll *b* and total chlorophyll) and leaf nitrogen content. Measurement of research variables for parameters is done by taking three plants per block per treatment which was indicated and measured at the end of harvest.

Plant wet weight was measured by taking the entire plant and weighed. Plant dry weight was measured by drying wet plant in an oven for 48 hours (until constant weight is obtained). Wet weight of fruit was obtained in every harvest (from harvest 1 to finish). Total weight of fruit is obtained by summing the wet weight of fruit at each harvest. Weight of fruit per plant was calculated by counting the total weight of the fruit divided by the total number of plants per plot (20 plants). Plant chlorophyll content was measured with a spectrophotometer according to Harborne (1987). Plant leaf nitrogen content was measured by using the method of Sudarmaji *et al.* (1996).

Data analysis

Data obtained from this study were analyzed by analysis of variance (ANOVA) and followed by DMRT (Duncan's multiple range test) using SPSS 10.05. In the analysis of variance, if *F* was greater than *F* table or a probability (sig) <0.05 hence *H*₀ refused and *H*₁ accepted.

DMRT (Duncan) populations that had the same average grouped into one subset. In one subset the treatment was not different (Hanafi 2005; Pratista 2002).

RESULTS AND DISCUSSION

Fertilization has long been known as a factor influencing the growth and yield of crops, including chili. Fertilizing is an effort to provide the necessary elements of plant nutrient. Provision of nutrients affects the levels of organic and inorganic compounds of the plant (Rosmarkam and Yuwono 2002). Among the parameters that can be observed as a physiological phenomenon is fresh weight, dry weight, chlorophyll and nitrogen content of leaves (Marschner 1986).

Biomass

Biomass in this study include the weight of wet and dry plants at the end of harvest and total wet weight of fruit per plant

Plant fresh weight

Fresh weight is the total weight of plants showing the results of metabolic activity. Fertilization can affect plant fresh weight as it provides nutrients from the soil. Fresh weight of pepper plants at different fertilizer treatment varied (Table 4). Fresh weight of the control group (P1) is equal to the weight of fresh manure treatment (P2), and not in one group with that treated with manure + chemical fertilizer + NPK (P3), and also that have different with that have treatment of chemical fertilizer + manure + liquid organic fertilizer (P4).

Table 1. Biomass plant two varieties of chili

Varietas	Biomassa			
	P1	P2	P3	P4
Plant fresh weight (g)				
<i>Fantastic</i>	228.93 a	496.01 ab	602.11 b	665.65 b
<i>Sakti</i>	277.16 c	508.33 cd	607.56 d	656.52 d
Plant dry weight (g)				
<i>Fantastic</i>	68.33b	111.11c	139.67cd	157 d
<i>Sakti</i>	126.59a	155.41a	354.28a	407.72a
Total fruit weight per plant (g)				
<i>Fantastic</i>	805.00 a	878.33 ab	770.83a	990.93 b
<i>Sakti</i>	530.83 c	546.67 c	606.67 cd	705.83 d

Note: Number that are noted same on the same line means that they are not significantly different according to DMRT test at *P* = 0.05. P1: Control, P2: Manure (2 kg/plant), P3: Manure (1 kg/plant) + Chemical fertilizer (ZA: KCl: SP-36 = 2: 1: 1) + NPK *Mutiara*, P4: manure + chemical fertilizer (KCl: SP-36 = 1: 1) + liquid organic fertilizer *Bathari Sri*.

Dry weight of plant

Dry weight of plant of both varieties of peppers varies. Various fertilization treatments significantly affect the increase in plant dry weight of *Fantastic* chili, but no significant effect on dry weight of *Sakti* chili (Table 5). The absence of influence of various fertilization treatments on

the dry weight of chili is a powerful indication that fertilization has not been quite able to meet the nutrient requirements needed by chili. Fertilizer treatment on different varieties of chili show different effect. This is in line with the statement Kartasapoetra (1995) and Abdul Rahim and Jumiati (2007) that the plant will need a variety of nutrients for growth and development, it takes them different time.

Although the fertilizer treatment was not significantly different in the dry weight of *Sakti*, but it can be seen that there is a tendency that fertilizer treatment can improve the dry weight of both varieties of chili. This means there is a tendency of photosynthesis to increase, due to the addition of nutrients from the soil as a result of the fertilization process.

Total fruit weight per plant

Total fruit weight per plant from both varieties of chili is different in various fertilizer treatments. The results showed that the fertilization treatment did not significantly affect fruit weight per plant in *Sakti* (Table 1). This gives an indication that the use of manure, chemical fertilizer and NPK (P3) is not always a solution to improve the yields. Only the precise formulation that increases the weight of pepper. Meanwhile, fertilizer treatment significantly affected total fruit weight per plant in *Fantastic* (Table 1). The presence of N, P, K fertilizer and organic matter deriving from chemical fertilizer and liquid organic fertilizer, make it possible to increase the weight of the chili. The combination of manure, chemical fertilizer and liquid organic fertilizer (P4) can be used to replace chemical fertilizer formulations that have been used by farmers in Klaten district.

Meanwhile in manure (1 kg/plant) + chemical fertilizer (ZA: KCl: SP-36 = 2: 1: 1) + Shake *Mutiara* NPK gave fruit weight per plant that was lower than the other treatments (Table 1). This can happen because the nitrogen in the treatment has exceeded the optimal point, thus causing partial breakaway of assimilated nitrogen as an amide, it just raises the nitrogen content of plants, and reduces the synthesis of carbohydrates (Rosmarkam and Yuwono 2005). Therefore, fresh weight of fruit that is formed is relatively lower than other treatments.

The increase in biomass of *Sakti* chili and the tendency of *Fantastic* chili biomass to increase show that the elements provided through fertilization function properly. Results of correlation analysis shows that fresh weight is positively associated with plant dry weight, chlorophyll *a*, chlorophyll *b* and total chlorophyll of the two varieties of the chili. This means that if the fresh weight of plants increases then the dry weight will increase too, and so are the chlorophyll *a*, chlorophyll *b* and total chlorophyll leaves of both varieties of chili.

Organic fertilizers can provide soil organic matter that is very helpful in restoring fertility of physics, chemistry and biology of soil, because it is useful to bind soil particles through the process of soil aggregation. Aggregation of soil can produce micro-pore space, so that the aeration in the soil becomes better and it creates optimum conditions for absorption of nutrients for plants

(Brady 1990). Influence of organic matter on soil chemical fertility, among others are cation and anion exchange capacity, the increase of soil microbial activity through decomposition and mineralization of organic matter (Suntoro 2002). Besides, organic material is also able to absorb and hold water (Juan et al. 2003), which in turn affects the accumulation of nutrients and products of metabolism that are stored in the fruit and seeds.

Meanwhile the chemical fertilizers progressively increase and complete nutrients (nitrogen, phosphorus, potassium, magnesium, sulfur) which are useful in increasing chili plant biomass. The nitrogen (from ZA and NPK *Mutiara*) is capable of acting as a constituent of many essential compounds such as proteins, amino acids, amides, nucleic acids, nucleotides, coenzymes, and many compounds essential to metabolism, a constituent of chlorophyll, cytosine and auxine hormones and as main components of plant dry matter. Nitrogen will increase the green color of leaves, stem and leaf growth (Marschner 1986). Nitrogen is closely related to the synthesis of chlorophyll (Sallisbury and Ross 1992) and the synthesis of proteins and enzymes (Schaffer 1996). Rubisco enzyme acts as a catalyst in the fixation of CO₂ that plants need for photosynthesis (Salisbury and Ross 1992; Schaffer 1996). Therefore, increasing the nitrogen content of plants can affect the good photosynthesis through chlorophyll content and photosynthetic enzymes, thereby increasing the *fotosintat* (fresh weight, dry weight, and weight of chili pepper) is formed.

The phosphorus (from SP-36) is an important component of ATP constituent compounds that act as an energy source in the dark reactions of photosynthesis and the nucleoprotein, the genetic information system (DNA and RNA), cell membranes (phospholipids), and phosphoprotein. KCl fertilization increases the availability and the absorbance of kalium, while the function of kalium in the chloroplast is to play role as a guard to keep a high pH. Kalium plays an important role in photosynthesis because it directly increases the growth and leaf area index, thus increases the assimilation of CO₂ and increases translocation and assimilation of photosynthesis result (Suntoro 2002).

Sulphur (from SP-36) is needed by plants to form the amino acids cystine, cysteine and methionine. Besides, sulfur is also part of biotin, thiamine, co-enzyme A and glutathione (Marschner 1995). Sulfur also functions as an activator, cofactor or regulator of enzymes and plays a role in the process of plant physiology. Elemental sulfur is an important part of peroxidase, an iron and sulfur compound contained in the chloroplast and is involved in oxydoreduction reaction with electron transfer and also in the reduction of nitrate in the process of photosynthesis (Tisdalle et al. 1990).

Dolomite can increase chlorophyll because the supply of Mg from dolomite are able to increase the availability of soil Mg and Mg plant uptake (Suntoro 2002). Artificial chemical fertilizers supplies certain nutrients in the form of highly concentrated inorganic compounds and easy to dissolve. Giving it repeatedly to plant can endanger the natural soil flora and fauna, bring in soil nutrient

imbalances, cause water pollution, especially ground water. Organic fertilizers supply various nutrients, especially in the form of a low concentration of organic compounds that are not easily soluble, so they do not cause nutrient imbalances in soil, even they can improve the nutrient balance. The supply of organic matter can nourish the life of the soil flora and fauna, which in turn will improve and maintain soil productivity (Nuryani and Sutanto 2002).

Significant increase in biomass, is a synergistic effect of organic extracts and chemical fertilizers in the soil. The supply of organic substrates through liquid organic fertilizer increases the activity of soil organisms that play a role in the decomposition of organic compounds and improve the physical fertility (soil structure, aggregation and aeration, increase water holding capacity) and increase nutrient availability (Reeves 1997). In such conditions the supply of oxygen which is required for respiration of soil organisms is available. Therefore, improvement of soil quality occurs in some phases and gives impact toward the increase *Sakti* chili biomass and tends to increase *Fantastic* chili biomass.

Chlorophyll content

Chlorophyll *a*

Fertilizer treatment gives effect on chlorophyll *a* content of two varieties of peppers (Table 2). Nutrients (nitrogen, phosphorus, magnesium, iron, manganese, potassium, calcium, sulfur) that accumulate in chemical fertilizer and organic fertilizer added to the manure treatment (1 kg/plant) + chemical fertilizer (ZA, SP-36, KCl = 2: 1: 1) + NPK *Mutiara* and manure (1 kg/plant) + chemical fertilizer (SP-36: KCl = 1:1) + liquid organic fertilizer can significantly increase the content of chlorophyll *a* in both varieties of chili. Chlorophyll *a* is in the leaves of both varieties of peppers varies. This is an indicator that the physiological response of these two varieties of chili against a given nutrient supply is different. In general, it is said that the supply of nutrients from fertilizer can increase chlorophyll *a* and both varieties of chili.

Table 2. Chlorophyll *a*, chlorophyll *b*, total chlorophyll at different fertilizer treatment given to two varieties of chili.

Varietas	Kandungan klorofil (mg/L)			
	P1	P2	P3	P4
Chlorophyll <i>a</i> (mg/L)				
<i>Fantastic</i>	2,52 a	5,99 ab	6,99 b	6,45 b
<i>Sakti</i>	2,65 a	6,29 b	7,48 bc	7,93 c
Chlorophyll <i>b</i> (mg/L)				
<i>Fantastic</i>	1,66 a	2,80 a	3,75 a	3,80 a
<i>Sakti</i>	1,68 a	3,23 ab	5,38 b	3,84 ab
Total chlorophyll (mg/L)				
<i>Fantastic</i>	4,17 a	8,78 ab	10,72 b	10,25 b
<i>Sakti</i>	4,33 a	9,51 b	13,31 b	11,31 b

Note: letters which are noted same on the same line means there were not significantly different according to DMRT test at P = 0.05. P1: Control, P2: Manure (2kg/tanaman), P3: Manure (1 kg/plant) Chemical Fertilizers (ZA: KCl: Sp-36 = 2: 1: 1) NPK *Mutiara*, P4: Fertilizer cage of chemical fertilizer (KCl: SP-36 = 1: 1) liquid organic fertilizer *Bathari Sri*

Chlorophyll *b*

Chlorophyll *b* functions as an antenna that gathers light and then transfer it to the reaction center. Reaction center is composed of chlorophyll *a*. Light energy is converted into chemical energy in the reaction center which can then be used for the reduction process in photosynthesis (Taiz and Zeiger 1991). Fertilizer treatment does not significantly affect chlorophyll *b* in two varieties of peppers (Table 2). This likely occurs because most of the chlorophyll is still at the stage of chlorophyll *a* (chlorophyll *a* of both varieties are proven to be significantly influenced by different fertilizer treatment) and has not become chlorophyll *b*, as we know that chlorophyll *a* is the precursor of chlorophyll *b* (Robinson 1980). Although fertilization treatments do not significantly affect the chlorophyll *b* content in both varieties of chili, but there is a tendency that fertilization can increase the content of chlorophyll *b* both varieties of chili.

Total chlorophyll

The use of manure (1 kg/plant) + chemical fertilizer (ZA, SP-36, KCl = 2: 1: 1) NPK *Mutiara* shows the highest total chlorophyll content for the two varieties of chili, because the artificial chemical fertilizers supply particular nutrients that are highly concentrated and easily soluble (N, P, K, Fe, Mg, S) that play a role in the formation of chlorophyll (Nuryani and Sutanto 2002). The control group had the lowest total chlorophyll content for both treatments, because there was no addition of nutrients from the outside, while the available nutrients in the soil were absorbed by the plant during the vegetative phase and early generative phase. Because of the low availability and low nutrient absorption then the formation of chlorophyll was disturbed. Therefore chili chlorophyll content of control group was relatively lower compared with other treatments.

The treatment of animal manure (1 kg/plant) + chemical fertilizer (SP-36: KCl = 1:1) liquid organic fertilizer is in the same group with the use of manure (1 kg/plant) of chemical fertilizer (ZA, SP-36, KCl = 2: 1: 1) NPK *Mutiara*. Nutrients (nitrogen, magnesium, iron, manganese), which accumulates in the chemical fertilizer that is added through fertilization can increase the total chlorophyll content of leaves of *Sakti* chili (Table 2). This condition is possible to happen because the element of N, P, K, Mg, Fe, and S which are chlorophyll-forming element are available and can be absorbed by plants.

After the analysis of variant it is known that fertilizer treatment doesn't significantly give effect on total chlorophyll of *Fantastic* chili, but significant effect on total chlorophyll of *Sakti* chili (Table 2). The existence of significant effect of fertilization on chlorophyll content indicates that the supply of nutrients (N, P, K, Mg, S) has a positive contribution to the process of leaf formation in *Sakti* chili. Meanwhile, *Fantastic* chili nutrient supply provided through fertilization has not been able to increase total leaf chlorophyll. This phenomenon indicates that the physiological response of both toward the fertilization is not the same, but there is a tendency that fertilization can increase the total chlorophyll content of leaves, and the manure and chemical fertilizer treatment (ZA: KCl: SP-36

= 2: 1: 1) NPK *Mutiara*) is always higher than any other treatments (Table 2). The full content of nutrients in the fertilizer formulations is able to provide a stimulant for the increase in total chlorophyll content for both varieties of chili.

The addition of organic matter increases the leaf chlorophyll, and the increase will be higher if it is accompanied by adding the dolomite and KCl. KCl fertilization increases the availability and uptake of phosphorus, while the function of potassium in the chloroplast acts as a guard to keep a high pH. Giving dolomite can increase chlorophyll because the supply of Mg from dolomite are able to increase the availability of soil Mg and Mg plant absorption. Magnesium plays a very important role in the synthesis of chlorophyll (Suntoro 2002; Rahayu 2002; Santi 2002). Giving ZA, NPK *Mutiara*, organic fertilizer can increase the chlorophyll because it is able to provide a combination of nitrogen and magnesium which are known as elements that absolutely must be available for the formation of chlorophyll (Dwijoseputro 1986).

Nitrogen is closely related to the synthesis of chlorophyll (Salisbury and Ross 1992) and synthesis of proteins and enzymes (Schaffer 1996). Rubisco enzyme acts as a catalyst in CO₂ fixation that plants need for photosynthesis (Salisbury and Ross 1992; Schaffer 1996). Therefore, total nitrogen content in plants can influence the outcome of photosynthesis via the photosynthetic enzymes and chlorophyll formation. In plants, nitrogen initially is in the form of ammonia and subsequent ammonia has been changed into glutamic acid, catalyzed by the enzyme glutamine synthetase (Harborne 1987). Glutamic acid serves as the base material in the biosynthesis of amino acids and nucleic acids (Nyakpa et al. 1988). Robinson (1980) called glutamic acid as a precursor for the porphyrin ring in formation of chlorophyll.

Chlorophyll formation mechanism begins with the formation of α aminolevulinic acid (ALA) (Stryer 2002). Formation of ALA through the stages of formation of glutamate through the glutamate-t-RNA, from glutamate and is converted into α ketoglut semialdehida then become the next-aldehyde with the enzyme transaminase or amino transferase enzyme ALA formed (Bonner and Varner 1965; Krogman 1979). Of the two molecules involves the enzyme ALA with dehydrated ALA will be formed porfobilinogen (PBG) containing pyrrole rings from four molecules of PBG to involve enzyme-ogen uroporfirin III. Decarboxylation changes uroporfirinogen III. Under aerobic conditions by involving kaproporfirinogen decarboxylase enzyme, kaproporfirinogen III will then be formed into proporfirinogen IX. Oxidation toward proporfirinogen IX will result in proporfirin IX which does not have Mg, after joining Mg then Mg protoporfirin IX is formed. The addition of methyl groups on protoporfirin IX Mg with the help of esterase protoporfirin will form Mg porphyrin IX monomethyl ester. Next is the change of Mg porphyrin IX monomethyl ester into pro-chlorophyllide (Bonner and Varner 1965; Devlin 1983; Krogman 1979).

Change from prochlorophyllide into chlorophyll takes place through the formation of proklorofilide holokrom

which binds to proteins that bind to ion 2H⁺, which will be given to the fourth ring which forms a holokrom proklorofilide, which can then be turned into chlorophyll *a* by releasing holokrom and apoprotein (Mohr and Schopfer 1995).

If chlorophyll *a* with the help of an enzyme that catalyzes the esterification compounds chlorophyllase phytol can be formed into chlorophyll *a*. Meanwhile leaf homogenate, and thylakoid supply and leaves that is protected from light can convert chlorophyll *a* to chlorophyll *b*. Therefore, chlorophyll *a* can become a precursor of chlorophyll *b* (Robinson 1980).

Results of correlation analysis shows that chlorophyll *a* has a positive relation with chlorophyll *b* and total leaf chlorophyll and is positively related to plant fresh weight. An increased in chlorophyll *a* will increase the chlorophyll *b*, total chlorophyll and fresh weight of plant leaves. This can be easily understood because chlorophyll *a* is a precursor for chlorophyll *b*, while the chlorophyll *a* and *b* is the composition of total leaf chlorophyll, and also part of the plant fresh weight.

Leaf nitrogen content

Results of the research shows that fertilizer treatments does not significantly influence the nitrogen content of leaves of two varieties of chili (Table 3). The results of this study supports the findings of the previous research which claims a combination of urea and organic fertilizer does not affect the nitrogen and chlorophyll content of hermada grass (Supriya and Soeharsono 2005). This is possible because in all treatment combinations, the plant minimum requirement for nitrogen has been met.

Therefore, although the nitrogen content provided through fertilization is high but the plants absorb only a certain number of plants as needed.

Table 3. Nitrogen content of leaves of two varieties of chillies on different fertilizer treatments.

Variety	Nitrogen content (mg/L)			
	P1	P2	P3	P4
<i>Fantastic</i>	3.64 a	3.46 a	3.85 a	4.12 a
<i>Sakti</i>	3.92 b	4.23 b	4.27 b	4.25 b

Note: letters which are same on the same line means not significantly different according to DMRT test at P = 0.05. P1: Control, P2: Manure (2 kg/plant), P3: Manure (1 kg/plant) + Chemical fertilizer (ZA: KCl: Sp-36 = 2: 1: 1) + NPK *Mutiara*, P4 : manure + chemical fertilizer (KCl: SP-36 = 1: 1) + liquid organic fertilizer

Although the nitrogen content of leaves at different fertilizer treatment is not significant, but there is a tendency that the fertilizer treatment can increase the nitrogen content of leaves (Table 3). The tendency of leaf nitrogen content increase is a reflection of the increased nitrogen that can be absorbed by plants. The combination of ZA fertilizer, NPK *Mutiara*, and organic fertilizers cause an increase in leaf nitrogen content of plants. Nitrogen that is available in the soil that can be absorbed by plant roots is in the form of nitrate ions and ammonium.

Both forms of nitrogen is obtained as a result of organic material decomposition. Nitrates that are absorbed by the roots and transferred into the upper parts of the plant is a result the leaf transpiration process. Thus, nitrate assimilation in higher plants generally occurs in the leaves.

The first step is the reduction of nitrate into ammonia. The second step, nitrite reaction which turn nitrite into nitrate that occurs to chlorophyll in the chloroplast. While the ammonia assimilation in most plants turn into glutamic acid. Glutamic acid serves as the base material in the biosynthesis of amino acids and nucleic acids (Harborne 1987; Nyakpa *et al.* 1988).

The tendency of increase in nitrogen content of plants can affect the photosynthesis either through chlorophyll content or photosynthetic enzyme. If the leaf nitrogen content increases, the fotosintat will increase, otherwise if the leaf nitrogen content decrease then fotosintat formed is also low. That's because the elements of nitrogen will increase the green color of leaves, support stem and leaf growth (Marschner 1986).

Nitrogen is closely related to the synthesis of chlorophyll (Sallisbury and Ross 1992) and synthesis of proteins and enzymes (Schaffer 1996). Rubisco enzyme acts as a catalyst in the fixation of CO₂ that plants need for photosynthesis (Salisbury and Ross 1992; Schaffer 1996).

Meanwhile, *Fantastic* chili leaf nitrogen content, manure treatment (1 kg/plant) + chemical fertilizer (SP-36: KCl = 1:1) + liquid organic fertilizer of 4.12%, followed by the use of manure + chemical fertilizer (ZA, SP-36, KCl = 2: 1: 1) + NPK *Mutiara* (3.85%), followed by the control group (3.64%) and the lowest manure application (2 kg/plant) amounted to 3.46% (Table 3).

The lower leaf nitrogen content in manure treatment than the control group at *Fantastic* chili, was caused by the nutrients in the manure treatment can not be absorbed optimally. This is related to the nature of the organic fertilizer that supplies a variety of nutrients, especially in the form of a low concentration of organic compounds that are not easily soluble (Nuryani and Sutanto 2002). The results of the research are consistent with research conducted by Salim (2006) which states that the type of soil management and organic fertilizer can not increase the total nitrogen, K-leaf and leaf nitrogen uptake.

CONCLUSION AND RECOMENDATION

In *Fantastic* chili, a variety of fertilization treatment affects the biomass and chlorophyll *a*, but it does not affect chlorophyll *b*, total chlorophyll and plant leaf nitrogen. The treatment of various kinds of fertilizers to *Sakti* chili effects on plant fresh weight, chlorophyll *a* and total chlorophyll and has no effect on dry weight, weight of fresh fruit, chlorophyll *b* and the nitrogen content of leaves. Researchers recommend to use the formulation of fertilizers with manure + chemical fertilizer (SP-36: KCl = 1: 1) + liquid organic fertilizer as a new alternative to chili cultivation which is more economical and environmental friendly.

REFERENCES

- Abdulrahim, Jumiaty. 2007. Effect of concentration and spraying time of liquid organic fertilizer super ACI on growth and yield of sweet corn 26 (3): 105-109. [Indonesia]
- Allabi DA. 2006. Effect of fertilizer phosphorus and poultry droppings treatments on growth and nutrient components of pepper (*Capsicum annuum* L.) African J Biotech 5 (8): 671-677.
- Bonner J, Varner JE. 1976. Plant biochemistry. 3rd ed. Academic Press. New York.
- Brady NC. 1990. The nature and properties of soils. 10th ed. Macmillan. New York.
- Chellemi DO, Lazarovits G. 2002. Effect of organic fertilizer applications on growth yield and pests of vegetable crops. Proc Fla State Hort Soc 115: 315-321.
- Devlin RM. 1983. Plant physiology. 3rd ed. D. Van Nostrand. New York.
- Diver S. 2001. Compost teas for plant disease control. ATTRA, Fayetteville, AR. <http://attra.ncat.org/attra-pub/compost-tea-notes.html>
- Dwijoseputro G. 1994. Introduction to plant physiology. Gramedia. Jakarta. [Indonesia]
- Foth HD. 1990. Fundamentals of soil science. 8th ed. John Wiley & Sons. New York.
- Gardner FP, Pearch RB, Mitchell RL. 1991. Physiology of crop plants. UI Press. Jakarta. [Indonesia]
- Guertal EA. 2000. Preplant slow release nitrogen fertilizers produce similar bell pepper yields as split applications of soluble fertilizers. Agronom J 92: 388-393
- Hanafiah KA. 2005. The experimental design. Raja Grafindo Persada. Jakarta. [Indonesia]
- Harborne JB. 1987. Phytochemical methods, guiding the modern way to analyze plant. Penerbit ITB. Bandung. [Indonesia]
- Juanda JSD, Assa'ad N, Warsana. 2003. Study on infiltration rate and some physical properties of soil on three types of hedgerows in alley cropping systems. J Ilmu Tanah dan Lingkungan 4 (1): 25-31. [Indonesia]
- Kartasapoetra AG. 1995. Climatology (The influence of climate on soil and plants). Bumi Aksara. Jakarta. [Indonesia]
- Kononova MM. 1999. Soil organic matter, its role in soil formation and soil fertility. Pergamon. London.
- Krogman DW. 1979. The biochemistry of green plant. Prentice Hall. New Delhi.
- Lakitan B. 2007. Fundamentals of plant physiology. Raja Grafindo Persada. Jakarta. [Indonesia]
- Loveless AR. 1991. Principles of plant biology to the tropics. Gramedia. Jakarta. [Indonesia]
- Marschner H. 1986. Mineral nutrition of higher plants. Academic Press. London.
- Mitchel P. 1994. Ecological methods for field and laboratory investigations. UI Press. Jakarta. [Indonesia]
- Mohr H, Schopfer P. 1995. Plant physiology. Springer. Berlin.
- Nelson DL, Cox MM. 2004. Lehninger principles of biochemistry. 4th ed. W. H. Freeman. New York.
- Nuryani dan Sutanto. 2002. Effect of municipal waste on the yield and nutrient captivity of chili. J Ilmu Tanah dan Lingkungan 3 (1): 24-28. [Indonesia]
- Nyakpa, Yusuf, Lubis AM, Pulung MA, Amran G, Munawar A, Go BH. 1988. Soil fertility. University of Lampung. Bandar Lampung. [Indonesia]
- Pracaya. 2000. Planting chili. Kanisius. Yogyakarta. [Indonesia]
- Prawirohatmodjo S, Marsoem SN, Sutjipto AH (eds). 2001. Environment conservation through efficiency utilization of forest biomass. Debut Press cooperation with Department of Forest Products Technology, Faculty of Forestry, Gadjah Mada University and JIFPRO (Japan International Forestry Promotion and Cooperation Center). Yogyakarta. [Indonesia]
- Rahayu H. 2002. Effect of additional doses of organic material and dolomite on the availability and uptake of P using peanut [*Arachis hypogaea* L. (Merr)] as indicator on the ground latosol. Sains Tanah 2 (1): 25-34. [Indonesia]
- Rauf A. 1995. Contribution of livestock waste in chili agribusiness in South Sulawesi. J Ilmiah Penelitian Ternak Gowa. Edisi Khusus. Sub Balai Penelitian Ternak Gowa. [Indonesia]
- Reeves DW. 1997. The role of soil organic matter in maintaining soil quality in continuous cropping systems. Soil Till Res 43: 131-167.

- Robinson T. 1980. The organic constituents of higher plants. 4th ed. Cordus Press. North Amherst, Mass. [Indonesia]
- Rosmarkam A, Yuwono NW. 2002. Soil fertility science. Kanisius. Yogyakarta. [Indonesia]
- Sadewa. 2008. Study of morphological and physiological growth of the vegetative phase of three varieties of chili pepper (*Capsicum annuum* L.) caused by the type of fertilizer. Faculty of Agriculture, University of Jember. Jember. [Indonesia]
- Salim AA. 2006. Effect of soil management and organic fertilizer on soil chemical properties, N uptake on leaves and yield of tea on andisols. J Penelitian Teh dan Kina 9 (1-2): 1-6. [Indonesia]
- Salisbury F, Ross CW. 2001. Plant physiology. Vol. 1, 2 and 3. ITB. Bandung [Indonesia]
- Santi. 2002. The effect of magnesium and cow dung extract on growth and yield of chili pepper (*Capsicum annuum* L.). University of Muhammadiyah Malang. Malang. [Indonesia]
- Schaffer AA. 1996. Photoassimilate distribution in plant and crops. Marcel Dekker. New York.
- Scheuerell SJ. 2004. Compost tea production practices, microbial properties, and plant disease suppression. International Conference on Soil and Compost Eco-Biology, September 15th – 17th 2004, León – Spain.
- Setiadi. 1993. Planting chili. Penebar Swadaya. Jakarta. [Indonesia]
- Simarmata T. 1999. Application of liquid organic fertilizer super bionic to increase fertilizer efficiency and land production towards sustainable agriculture. PT. Foreverindo Insan Abadi. Jakarta. [Indonesia]
- Simarmata T. 2005. Application of organic extracts to improve the efficiency of chicken manure on Inceptisols with tomato yield as indicators. Agrikultura 16 (2): 84-88. [Indonesia]
- Stryer L, Berg JM, Tymoczko JL. 2002. Biochemistry. 5th ed. WH Freeman. San Francisco.
- Subagyo H. 1970. Principles of soil science. Soeroengan. Jakarta [Indonesia]
- Sudarmadji S, Haryono B, Suhardi. 1996. Analysis of food and agriculture. Liberty. Yogyakarta. [Indonesia]
- Suntoro. 2002. Effect of organic matter addition, dolomite and KCl on chlorophyll content and its impact on peanut (*Arachis hypogaea* L.) yield. BioSMART 4 (2): 36-40. [Indonesia]
- Supriadi, Soeharsono. 2005. The combination of urea and organic fertilizers on inceptisol soil to physiological response of hermada grass (*Sorghum bicolor*). Proceedings of the National Seminar on Technology of Animal Husbandry and Veterinary Science, Bogor, 12-13 September 2005. [Indonesia]
- Sutejo MM. 2002. Fertilizer and how fertilization. Rineka Cipta. Jakarta. [Indonesia]
- Taiz L, Zeiger E. 1991. Plant physiology. Benyamin/Cumming. Tokyo.
- Tandisau P, Sariubang M. 1995. Manure and its relationship with soil fertility and cotton yield. J Ilmiah Penelitian Ternak Gowa. Edisi Khusus. 1-6. [Indonesia]
- Tisdale SL, Nelson WL, Beaton JD. 1985. Soil fertility and fertilizer. Macmillan. New York.